



American College of
Emergency Physicians®

ADVANCING EMERGENCY CARE 

POLICY STATEMENT

Approved September
2018

Pediatric Mental Health Emergencies in the Emergency Department

Revised September 2018 with
current title

Reaffirmed April 2012

Originally approved April 2006
titled "Pediatric Mental Health
Emergencies in the Emergency
Medical Services System"

ABSTRACT

Emergency departments (EDs) are vital in the management of pediatric patients with mental health emergencies (MHE). Pediatric MHE are an increasing part of emergency medical practice. In their capacity as a safety net resource for many healthcare problems, EDs are increasingly used for mental health visits. A fragmented mental health infrastructure that is critically low in resources and staffing within all sectors significantly contributes to this problem. EDs are tasked to safely and humanely manage children with known mental illness and those with an acute behavioral crisis while utilizing a culturally and developmentally appropriate approach. The ED is tasked to provide initial management and stabilization of patients within the entire spectrum of mental health, from suicidality, maltreatment, and post-traumatic stress to violence, aggression, autistic spectrum disorder, and substance abuse. It is critical that ED's address not only the physical but also the mental health needs of patients during and after mass casualty incidents and disasters. Yet, many ED providers often feel that they don't have the required training and expertise to serve beyond that of a frontline resource or crisis management for mental health emergencies. Robust coordination with the mental health community is therefore essential for the management of these patients across the continuum from crisis management to ongoing care.

The American College of Emergency Physicians supports the following actions: advocacy for increased community mental health resources and linking them to the medical home, EDs, and inpatient psychiatric hospitals, as well as improved pediatric mental health tools for the ED, increased mental health insurance coverage, adequate reimbursement at all levels; acknowledgment of the importance of the child's medical home and their role in managing crisis events, development of community paramedicine programs for accurate assessment and triage of behavioral health crisis, and promotion of education and research for mental health emergencies.

Key words: Emergency department, mental health emergencies, school and community mental health services, medical home

INTRODUCTION

Pediatric mental health emergencies constitute a large and growing segment of pediatric emergency department (ED) volume. The ED, therefore, plays a critical role in the evaluation and management of children and adolescent patients with mental health emergencies. Community mental health resources have become diminished and, in some regions, disappeared entirely through inpatient bed shortages, private and public health insurance changes, reorganization of state mental health programs,¹ and shortages of providers who specialize in pediatric mental health.² These changes have resulted in critical shortages for both inpatient and outpatient mental health services for children.³ The ED has increasingly become the safety net for a fragmented mental health infrastructure in which the needs of children and adolescents, among the most vulnerable populations, have not been sufficiently addressed.

EDs should safely, humanely, and in a culturally sensitive manner manage patients with exacerbations of known diagnosed mental illnesses as well as those with developmental delay, autistic spectrum disorders, ADHD, or those in behavioral crisis. EDs also should identify and manage patients with previously undiagnosed, undetected or emerging mental health related conditions such as suicidal ideation, depression, escalating aggression, substance abuse, and post-traumatic stress disorder. Given the broad scope of mental health emergencies, EDs need to also consider and be able to treat mental health manifestations of trauma, physical and sexual maltreatment, and in children exposed to community and domestic violence. Violence-related situations may involve pediatric victims and/or pediatric-aged perpetrators of violence. In many states, adolescents can seek and receive care for mental health issues and drug/alcohol use without parental involvement, and EDs should maintain confidentiality unless the child is at risk of harming himself/herself or others. The ED must also recognize the primary support role of the family and caregivers in all phases of pediatric mental illness.

EDs also play a critical role in mass casualty occurrences and disasters and must address the unique mental health needs of children during and after these events. A strong and growing body of evidence indicates that emotional and physical trauma to children can cause neurochemical and structural brain changes resulting in post-traumatic stress disorder, and this can affect some children into their adult lives.³⁻¹⁴ Emotional trauma may be ameliorated by timely, culturally-appropriate, pediatric-specific stress intervention that may be implemented in the initial hours after the trauma.^{15,16}

The epidemiologic and outcome data on pediatric mental health emergencies are insufficient, but there is evidence that pediatric mental health concerns are commonly unaddressed.^{17,18} Pediatric mental health emergencies are frequently not recognized as such, presenting initially as trauma or somatic complaints, and are, therefore, underrepresented in the existing data.¹⁹⁻²² The challenges to an already overburdened ED "safety net" are to provide safe, humane, and culturally and developmentally sensitive triage, diagnosis, stabilization, initial management, and treatment and referral for a broad spectrum of mental health emergencies, working within a mental health infrastructure in crisis.

Early identification and intervention can aid in preventing the emergency behavioral health crisis. This in turn has been shown in some community-based programs to effectively decrease admission rates and decrease the length of ED stays.²⁵ Some EMS agencies have developed community paramedicine programs to more accurately assess and triage behavioral health crises to affect optimal delivery of care.^{26,27} In this regard, the patient may access the medical home for direct behavioral health referral thus avoiding the emergency department as a needed intervention.²⁴ Such efforts would alleviate an already strained emergency system that currently experiences excess boarding rates for behavioral health patients where the skilled mental health team is not available.

Pediatric mental health emergencies are best managed by a skilled, multidisciplinary team approach, including specialized screening tools,²⁸ pediatric-trained mental health consultants, the availability of

pediatric psychiatric facilities when hospitalization is necessary, and an outpatient infrastructure that supports pediatric mental health care, including communication back to the primary care physician and timely and appropriate ED referrals to mental health professionals.²³

The American College of Emergency Physicians supports the following actions:

1. Advocacy for adequate pediatric mental health resources in both inpatient and outpatient settings, including the availability of prompt psychiatric consultation for ED psychiatric patients, as well as school and community mental health services, including adequate mental health screening.
2. Development of mechanisms for the ED to train all ED staff in dealing with unique pediatric mental health issues and to properly recognize and respond to cases, including violence in the community, physical trauma, domestic violence, child maltreatment, mass casualty incidents and disasters, suicides and suicide attempts.
3. Appropriate reimbursement for both inpatient and outpatient pediatric mental health services.
4. Development of cross institutional transfer agreements with specific mental health facilities to simplify the process of bed search and placement for patients requiring inpatient mental health care. Such efforts should be directed towards limiting emergency department length of stay and boarding.
5. Acknowledgment of the importance of the child's medical home to his or her continued well-being, including prevention, screening, crisis intervention, and treatment of mental health issues.²²
6. Advocacy for comprehensive insurance coverage for pediatric mental health to include provision of services for the uninsured, as well as expansion of coverage to include mental health services for those who are insured.
7. Advocate for community-based behavioral services using a patient centered approach to identify and manage behavioral health concerns prior to development of an emergent condition. This includes efforts directed towards ED screening to identify suicide risk and methods of suicide prevention.
8. Advocacy for additional research funding dedicated to pediatric emergency mental health issues.
9. Promotion of education and research for mental health emergencies, specifically:
 - To expand the data on epidemiology, best practices, treatment outcomes, and cost-effectiveness of management of pediatric mental health emergencies in the ED.
 - To evaluate the adequacy of patient access to pediatric mental health services.
 - To evaluate children with behavioral crisis and to understand gaps in primary care and community resources.
 - To develop mental health support networks that minimize reliance on acute crisis management.
 - To develop and validate accurate pediatric mental health screening tools for use in various settings and best practices for follow-up programs for pediatric mental health patients.
 - Acknowledge and support community-based integration of services utilizing tools such as telemedicine and paramedicine programs adequately trained to identify and assess crisis events for appropriate triage and transport to behavioral health crisis facilities as an alternative to emergency medical care facilities.
 - To enhance the pediatric mental health educational needs of practicing EM physicians and the curriculum for emergency medicine and pediatric residency training programs and pediatric emergency medicine fellowships.

An earlier version of this policy statement has been approved by the American College of Emergency Physicians Board of Directors and the American Academy of Pediatrics Board of Directors.²⁹

REFERENCES

1. Okumura MJ, Knauer HA, Calvin K, et al. Caring for children with special health care needs: profiling pediatricians and their health care resources. *Matern Child Health J.* 2018 Jul;22(7):1042-1050.

2. Leslie LK, Baum R, Turner A. Revisiting the viability of the developmental-behavioral health care workforce. *Pediatrics*. 2018 Mar;141(3):e20174132.
3. American Academy of Pediatrics Committee on Child Health Financing. Scope of health care benefits from birth through age 21. *Pediatrics*. 2006 Mar;117(3):979-982.
4. American Academy of Pediatrics Committee on Injury and Poison Prevention. Firearm-related injuries affecting the pediatric population. *Pediatrics*. 2000 Apr;105(4):888-895.
5. Terr LC. Childhood traumas: an outline and review. *Am J Psychiatry*. 1991 Jan;148(1):10-20.
6. Brick ND. The neurological basis for the theory of recovered memory [research paper]. AOL Hometown. 2003. Available at: http://members.aol.com/smartnews/Neurological_Memory.htm. Accessed August 7, 2005.
7. Meichenbaum D. *A Clinical Handbook/Practical Therapist Manual for Assessing and Treating Adults with Post-Traumatic Stress Disorder (PTSD)*. Waterloo, Ontario: Institute Press; 1994.
8. Foy DW. Introduction and description of the disorder. In: Foy DW, ed. *Treating PTSD: Cognitive-Behavioral Strategies*. New York, NY: Guilford; 1992:1-12.
9. Knopp FH, Benson AR. *A Primer on the Complexities of Traumatic Memory Childhood Sexual Abuse: A Psychobiological Approach*. Brandon, VT: Safer Society Press; 1996.
10. van der Kolk BA. *The Body Keeps the Score: Memory and the Evolving Psychobiology of Post Traumatic Stress*. 1994. Available at: <http://www.trauma-pages.com/vanderk4.htm>. Accessed August 7, 2005.
11. Foy D. *Scientific American*. 1995;273(4):14.
12. Winston FK, Kassam-Adams N, Vivarelli-O'Neill C, et al. Acute stress disorder symptoms in children and their parents after pediatric traffic injury. *Pediatrics*. 2002 Jun;109(6):e90.
13. Children's Hospital of Philadelphia. Post-traumatic stress disorder may follow traffic crashes according to doctors at the Children's Hospital of Philadelphia [press release]. Available at: http://www.eurekalert.org/pub_releases/1999-12/CHoP-Psdm-061299.php. Accessed August 7, 2005.
14. American Academy of Pediatrics. Insurance coverage of mental health and substance abuse services for children and adolescents: a consensus statement. *Pediatrics*. 2000;106:860-862.
15. Davidhizar R, Shearer R. Helping children cope with public disasters. *Am J Nurs*. 2002;102:26-33.
16. Kalyjian A. Sri Lanka: post tsunami mental health outreach project—lessons learned. *Psi Chi: The National Honor Society in Psychology*. February 24, 2005.
17. Popovic JR. 1999 National Hospital Discharge Survey: annual summary with detailed diagnosis and procedure data. *Vital Health Stat 13*. 2001;13(151):i-v, 1-206.
18. Olson L, Melese-d'Hospital I, Cook L, et al. Mental health problems of children presenting to emergency departments. Paper presented at: Third National Congress on Childhood Emergencies; April 15-17, 2002; Dallas, TX.
19. US Consumer Product Safety Commission, Division of Hazard and Injury Data Systems. *Hospital-Based Pediatric Emergency Resource Survey*. Bethesda, MD: US Consumer Product Safety Commission; 1997.
20. Seidel JS, Hornbein M, Yoshiyama K, et al. Emergency medical services and the pediatric patient: are the needs being met? *Pediatrics*. 1984;73:769-772.
21. Seidel JS. Emergency medical services and the adolescent patient. *J Adolesc Health*. 1991;12:95-100
22. Sapien RE, Fullerton L, Olson LM, et al. Disturbing trends: the epidemiology of pediatric emergency medical services use. *Acad Emerg Med*. 1999;6:232-238.
23. American Academy of Pediatrics. Indications for management and referral of patients involved in substance abuse. Committee on Substance Abuse. *Pediatrics*. 2000;106:143-148.
24. American Academy of Pediatrics, Medical Home Initiatives for Children with Special Needs Project Advisory Committee. The medical home. *Pediatrics*. 2002;110:184-186.
25. Sheridan DC, Sheridan J, Johnson KP et.al. The effect of a dedicated psychiatric team to pediatric emergency mental health care. *J Emerg Med*. 2016 50, e121-8.
26. Patterson DG, Coulthard C, Garberson LA, et al. What is the potential of community paramedicine to fill rural health care gaps? *J Health Care Poor Underserved*. 2016;27(4A):144-158.

-
27. Choi BY1, Blumberg C2, Williams K3. Mobile integrated health care and community paramedicine: An emerging emergency medical services concept. *Ann Emerg Med.* 2016 Mar;67(3):361-6.
 28. King CA, O'Mara RM, Hayward CN, et.al. Adolescent suicide risk screening in the emergency department. *Acad Emerg. Med.* 2009;16:1234-1241.
 29. American College of Emergency Physicians. Pediatric Mental Health Emergencies in the Emergency Medical Services System. *Ann Emerg Med.* 2006 Oct;48(4):484-6.