

PO BOX 649 FORT DEFIANCE, AZ 86504 PHONE: 928.729.8000 FAX: 928.729.8019 WEBSITE: <u>WWW.FDIHB.ORG</u>

ED CHECKLIST FOR SEPSIS - TO ORGANIZE AND ENHANCE BEDSIDE CARE

1.	First 15-30 minutes				
	☐ Screen patient for spot sepsis criteria, Physician decides to call				
	"Code Sepsis".				
	☐ Establish 2 large bore IV's, 30CC/kg initial fluid bolos over 30'				
	□ Sepsis order set/ Draw Labs/ Lactate/ Blood Culture				
	☐ Start antibiotics- use TMC Antimicrobial guidelines for suspected				
	source (see desktop share drive)				
	□ VS q 15' report to physician				
2.	30 minutes – 3 hours				
	 Re-evaluate perfusion and volume status 				
	☐ Continue fluid resuscitation as appropriate (make sure patient fluid				
	responsive/not in ARDS)				
	☐ Recheck Lactate at 3 hours (goal >10% clearance)				
	□ Pressors if MAP <65 despite adequate volume (initiated pressor of				
	choice is norepi				
	□ Facilitate ICU transfer				
	□ Accomplish ancillary studies / review of lab for findings				