

EINSTEIN EMERGENCY MEDICINE

Adult Sepsis Bedside Tool

Date:

Patient Weight:

VITAL SIGNS: T

HR

RR

BP

SIRS SIGNS: T >38 or < 36

HR >90

RR > 20

WBC > 12, < 4 or > 10% bands

EARLY RECOGNITION: SIRS SIGNS AND SUSPECTED INFECTION

	TIME (24hr)	GOAL (cumulative time)
Triage	_____	<10 minutes
Recognition of Sepsis	_____	<10 minutes
Severe Sepsis?	YES?	NO?
ED Provider to Bedside	_____	<10 minutes

IV ACCESS AND FLUID BOLUS

IV Access	_____	<15 minutes
IVF BOLUS 30cc/kg Start time (Use Pressure bag)	_____	_____
Contraindications to full bolus?	_____	
IVF End time	_____	<1 hour

CULTURES and LABS

Labs drawn	_____	_____
Initial Lactate: _____	SEVERE SEPSIS?	_____
Blood cultures/urine cultures	_____	_____
Repeat lactate: _____	SEVERE SEPSIS?	_____

EARLY ANTIBIOTICS AND SHOCK MANAGEMENT

Antibiotics	_____	<3 hours
Need vasopressors?	YES/NO	

EARLY CRITICAL CARE CONSULT AND ED2 TIME

MICU or Surgery Paged	_____	_____
Admitting Consultant @ Bedside	_____	_____
Patient transported to ICU/OR/SDU	_____	_____